

Autonomous Mobile Robots

CHEAT SHEET BY JANIS HUTZ
ETHZ, FS2026

1 Introduction

1.1 Probability

D 1.1 (Sum rule) $P(X) = \sum P(X, Y) = \sum P(X \cap Y)$

D 1.2 (Prod) $P(X, Y) = P(X|Y)P(Y) = P(Y|X)P(X)$

T 1.3 (Bayes) $P(Y_i|X) = \frac{P(X|Y_i)P(Y_i)}{\sum_{j=1}^n P(X|Y_j)P(Y_j)}$

D 1.4 (Cont. Var) Sums become integrals
e.g. $\sum_X P(X) = 1$ becomes $\int p(x) dx = 1$

D 1.5 (Indep.) x, y indep. iff $p(x, y) = p(x)p(y)$

D 1.6 (Cond. Indep.) iff $p(x, y|z) = p(x|z)p(y|z)$

D 1.7 $E[x] = \int_{-\infty}^{\infty} xp(x) dx$, also for $x = f(x)$

D 1.8 $\text{Cov}[x] = E[xx^T] - E[x]E[x]^T = \Sigma$

D 1.9 (Gauss. Dist.) $x \sim \mathcal{N}(\mu, \Sigma)$ (μ mean, Σ cov.),
PDF: $p(x) = \frac{1}{\sqrt{(2\pi)^k |\Sigma|}} \exp(-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu))$

1.2 Measurement models

$z = b_C + sM_S\omega + b + n + o$: b_C const bias, b time bias, M missal., $n \sim \mathcal{N}(0, R)$ noise, $s\omega$ corr. meas., o other infl.

2 Locomotion & Kinematics

2.1 Positioning

D 2.1 (Position Vector) ${}^W t_B = {}^W t_W {}^W t_B$, Original Frame, End point, Target Frame, $\sin = s$, $\cos = c$

D 2.2 (State vector) x_R : x, v of rob in W , pos of sensors

D 2.3 (Rot. Mat.) $R_z = \begin{bmatrix} c(\psi) & -s(\psi) & 0 \\ s(\psi) & c(\psi) & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$$R_y(\theta) = \begin{bmatrix} c(\psi) & 0 & s(\psi) \\ 0 & 1 & 0 \\ -s(\psi) & 0 & c(\psi) \end{bmatrix} R_x(\varphi) \begin{bmatrix} 1 & 0 & 0 \\ 0 & c(\psi) & -s(\psi) \\ 0 & s(\psi) & c(\psi) \end{bmatrix}$$

R 2.4 Application: ${}_W a = R_{WB} a$

L 2.5 $R_{BW} = R_{WB}^{-1} = R_{WB}^T$, $\det(R_{WB}) = 1$ (orth.)

R 2.6 Cols of R_{WB} are basis vec. of Frame \vec{F}_B in \vec{F}_W

D 2.7 (Euler Angles) Yaw (z), Pitch (y), Roll (x), mult. rotation matrices, e.g. $R_{EB} = R_z(\psi) \cdot R_y(\theta) \cdot R_x(\varphi)$, bound. $[n]^\times = n x^T$ (matrix from vec + arg x)

D 2.8 (Rot. Vec) $\alpha = \alpha n$ (n normal)

$$R(\alpha, n) = I_3 + \sin(\alpha)[n]^\times + (1 - \cos(\alpha))([n]^\times)^2$$

D 2.9 (Quaternions) $q = q_w + q_x i + q_y j + q_z k$ with $i^2 = j^2 = k^2 = -1$, $(ij = -ji = k, \text{ same for } jk \text{ and } ki)$

D 2.10 (Transf. M) $T_{AB} = \begin{bmatrix} R_{AB} & A t_B \\ \mathbf{0}_{1 \times 3} & 1 \end{bmatrix}$

$$T_{BA} = T_{AB}^{-1} = \begin{bmatrix} R_{AB}^T & -R_{AB}^T A t_B \\ \mathbf{0}_{1 \times 3} & 1 \end{bmatrix} \quad T_{AC} = T_{AB} T_{BC}$$